

## Racumin® Tracking Powder

### Characteristics

- 1 Multiple feed anti-coagulant rodenticide
- 2 Unique formulation
- 3 Resistance management (overcomes bait shyness)
- 4 Cost effective
- 5 Contains a bittering agent that prevents non-target animals and humans from ingesting the product.

Pack size: 5 kg

### Product Name:



**Racumin® Tracking Powder** - REG. NO. L2800 Act No. 36 of 1947

An anti-coagulant poison for control of the Norway Rat, Roof Rat and House Mouse in and around human and animal dwellings, factories and warehouses.

UN NO. 3027

CAUTION

### Active Ingredient

- 1 *Coumatetralyl (anti-coagulant) 7,5 g/kg*

### Warnings

- 1 Handle with care.
- 2 Poisonous when swallowed and inhaled.
- 3 Toxic to fish, wildlife, pigs and other domestic animals.
- 4 Store under lock and key away from food and feedstuffs.
- 5 Keep out of reach of children, animals and uninformed persons, poultry and livestock.
- 6 In case of poisoning CALL A DOCTOR AND MAKE THIS INSERT AVAILABLE TO HIM.

### Precautions

- 1 Do not inhale dust.
- 2 Wear rubber gloves when handling powder or bait.
- 3 Wash with soap and water immediately after use or accidental skin contact.
- 4 Do not eat, drink, or smoke whilst mixing or applying or before washing hands and face.
- 5 Clean or destroy all mixing utensils and dispose of wash water where it will not contaminate crops, grazing, rivers, dams.
- 6 Destroy empty container and never use for any other purpose.
- 7 Prevent contamination of any edible commodity, drinking water, eating utensils and areas not under treatment.

- 8 The bait or powder should be set out in such places where it will be completely inaccessible to children and domestic animals. The use of bait boxes or other special containers is strongly recommended.
- 9 Unconsumed bait residues must be removed and burned or buried at the end of treatment.

### First Aid

- 1 Skin contact – immediately remove contaminated clothing. Wash affected area with plenty of water and soap for at least 20 minutes.
- 2 If accidentally swallowed induce vomiting by tickling the back of the throat. Repeat until vomit is clear.

### Directions for Use

- 1 Use only as directed.

### Pest

#### Dosage

#### Application directions

Norway

Rat,

Roof Rat

Undiluted

#### Tracking Powder:

The **Racumin® Tracking Powder** should be sprinkled, not too thinly, in rat holes, on rat runs and around hiding places. Leave the powder down for at least 5 days and replenish as required.

One part +  
15-20 parts  
bait material

#### Bait:

Thoroughly mix one part by mass of **Racumin® Tracking Powder** with 15-20 parts of bait material e.g. 30 g in 450-600 g bait material and set out in places frequented by rats. Very good effect is obtained with dry baiting material made up as follows: 60 % crushed maize, 35 % rolled oats, 5 % sugar. Traces of finely cut raw bacon. The bait must be laid out on at least 5 successive days or as long as the rats consume it. It is essential that the bait remains in a good, attractive condition during this period.

House

Mouse

As above

Apply the powder and poisoned baiting material in the same way as for rats.

**Racumin®** is strongly recommended for joint control of mice and rats or when it is uncertain from damage caused whether the area is infested by rats or mice.

#### NOTE:

- 1 Coumatetralyl, like other coumarin compounds has a cumulative action i.e. small quantities must be consumed on several consecutive days until the specific symptoms of poisoning appear.
- 2 Consequently, it is the repeated intake of small dosages and not a single intake of a large quantity that causes eradication of the rodents.
- 3 This multiple feed efficacy reduces the risk of secondary poisoning of non-target animals.

#### Note Physicians



- 1 Intravenous Vitamin K1, 50 - 150 mg on first day depending on prothrombin time.
- 2 In severe cases small transfusions of fresh blood.

### Symptoms

- 1 Bleeding from the gums and nose, bleeding under the skin, pallor, haemorrhage into and around elbow and knee joints.
- 2 Blood in urine and faeces.

